

Random Quantum Spin Chains: A Real-Space Renormalization Group Study

E. Westerberg, A. Furusaki,* M. Sigrist, and P. A. Lee

Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Quantum Heisenberg spin chains with random couplings and spin sizes are studied using a real-space renormalization group technique. These systems belong to a new universality class of disordered quantum spin systems realized in *e.g.* $\text{Sr}_3\text{CuPt}_{1-x}\text{Ir}_x\text{O}_6$. The low-energy fixed point is characterized by the formation of weakly coupled large effective spins. At low temperature T the entropy obeys a power law T^δ ($\delta \approx 0.44$), and the susceptibility follows a Curie-like behavior.

A class of quasi-one-dimensional (1D) spin systems with an interesting type of disorder was recently discovered in the compound $\text{Sr}_3\text{CuPt}_{1-x}\text{Ir}_x\text{O}_6$. In the pure material, the Cu (spin $S = \frac{1}{2}$) and Pt ($S = 0$) alternate along chains and the Cu spins interact antiferromagnetically with each other [1]. In the compound $\text{Sr}_3\text{CuPt}_{1-x}\text{Ir}_x\text{O}_6$ a fraction of the spinless Pt atoms is replaced by Ir with a spin $S = \frac{1}{2}$ that couples ferromagnetically to the neighboring Cu spins. Thus this alloy represents a quasi-1D system with random ferromagnetic (FM) and antiferromagnetic (AF) bonds of fixed strength [2]. Measurements of the uniform susceptibility as a function of temperature ($2\text{K} < T < 300\text{K}$) revealed a crossover around 10K between the standard high-temperature and a low-temperature Curie behavior [3].

In a recent theoretical study we considered a related model of a quantum spin chain with random FM and AF nearest-neighbor coupling of equal strength J [4]. The analysis of high-temperature series for the susceptibility compares qualitatively well with the experimental data [5]. Our analysis of the entropy led to the surprising result that a considerable fraction of the degrees of freedom remains uncorrelated down to a temperature $k_B T \sim J/5$. This unusual property together with the Curie-like behavior of the susceptibility indicated the existence of very weakly coupled effective spin degrees of freedom in the low-energy regime, which we identify in the following way. Our system consists of a sequence of alternating FM and AF segments of random length. At an intermediate energy scale ($\sim J$) the spins within each segment lock into their local ground state with a variety of spin values corresponding to the smallest or largest possible total spin for AF or FM segments, respectively. Because of the quantum nature of the spins these effective degrees of freedom couple rather weakly and behave like independent spins for a certain range of temperature. They provide the effective spins seen in the Curie behavior below the crossover and account for the entropy below $k_B T \sim J/5$ [4]. At a very low energy scale these effective spins will correlate among each other. This picture has been confirmed by numerical studies of finite size chains which show that the low-energy dynamics is

described by effective spins of varying sizes coupled by an exchange interaction whose strength and sign depend on the length and order of the segments [6]. This suggests that in order to study the low-temperature properties of the random spin- $\frac{1}{2}$ chain, it is necessary to introduce an effective Hamiltonian,

$$\mathcal{H} = \sum_i J_i \mathbf{S}_i \cdot \mathbf{S}_{i+1}, \quad (1)$$

where the strength and sign of J_i as well as the size of the spins S_i are random. The great generality of the Hamiltonian (1) suggests that apart from describing the low-energy behavior of $\text{Sr}_3\text{CuPt}_{1-x}\text{Ir}_x\text{O}_6$ it should be applicable to a wider range of disordered 1D spin systems.

We investigate the low-temperature behavior of systems described by (1) by means of a generalized version of the real-space renormalization group (RG) method invented by Dasgupta and Ma (DM) [7] to study random bond AF $S = \frac{1}{2}$ Heisenberg chains (see also [8,9]). Let us define a link in the chain as two neighboring spins and the coupling J connecting them. If a link were isolated from the rest of the chain, it would form a local ground state of maximum ($J < 0$) or minimum ($J > 0$) spin with an energy gap Δ to the first excited multiplet. For a FM bond $\Delta = -J(S_L + S_R)$ while for an AF bond $\Delta = J(|S_L - S_R| + 1)$, where S_L and S_R are the left and right spins of the link, respectively. If we keep track of whether a link is ferromagnetic or antiferromagnetic, there is a one-to-one correspondence between gaps and coupling constants and we characterize from now on a link by $\{\Delta, S_L, S_R\}$. Now we focus on the strongest link in the chain, defined as the link with the largest gap $\Delta \equiv \Delta_0$. If the distribution of gaps is broad, the gaps of the two neighboring links, Δ_1 and Δ_2 , are typically much smaller than Δ_0 so that the two spins S_L and S_R , to a good approximation, lock into their local ground state. Consequently we replace the link $\{\Delta_0, S_L, S_R\}$ by a single effective spin $\tilde{S} = |S_L \pm S_R|$ representing the local ground state of minimum (AF) or maximum (FM) spin. The weaker neighboring bonds are then taken into account perturbatively in $\Delta_{1,2}/\Delta_0$, leading to an effective interaction between the spins S_1 , $\tilde{S}$, and S_2 (see inset in Fig. 1). Spin isotropy is conserved in this procedure, and

to first order in $\Delta_{1,2}/\Delta_0$ no next-nearest-neighbor interactions are generated. The case of $\tilde{S} = 0$ (corresponding to an AF link with $S_L = S_R$) needs a special treatment, since to first order this would simply decouple the left and right part of the chain. In this case we remove both S_L and S_R (which together form a singlet) and make use of the fact that second-order processes introduce a weak effective coupling between S_1 and S_2 . This last step corresponds to the DM treatment of the purely AF spin- $\frac{1}{2}$ model.

Integrating out the strongest link successively in the manner described above preserves the form of the Hamiltonian (1) but changes the distribution of links and, in particular, lowers Δ_0 . Thus the RG procedure generates a flow in the distributions of FM and AF links,

$$P^{F,A}(\Delta_0; \Delta, S_L, S_R). \quad (2)$$

After having integrated out links until the largest remaining gap in the chain is Δ_0 , the chain is described by the Hamiltonian (1) with AF and FM links distributed according to (2). For low enough energies Δ_0 we expect the distributions to flow to fixed-point distributions where P^F and P^A exhibit scaling behavior

$$P^{F,A}(\Delta_0; \Delta, S_L, S_R) = \Delta_0^{-\gamma} Q^{F,A}\left(\frac{\Delta}{\Delta_0^\beta}, \frac{S_L}{\Delta_0^{-\alpha}}, \frac{S_R}{\Delta_0^{-\alpha}}\right). \quad (3)$$

In principle the exponents α , β , and γ in P^F could differ from the ones in P^A . However, this is very unlikely, because FM and AF links are strongly interrelated with each other as we can see from the fact that a large number of the spins belong to both an AF and a FM link and that in the RG procedure an AF link can be converted into a FM link and vice versa. Moreover, different exponents γ would imply that one of the probability distributions scales to zero, leaving us with a purely FM or AF chain. Indeed our numerical RG calculation shows that this is not the case. In addition to the exponents in (3) we are interested in how length scales with Δ_0 . Defining n as the ratio of the original number of sites to the number of effective spins, we expect $n \sim \Delta_0^{-\delta}$.

The exponents in (3) are not all independent. From the normalization condition $\sum_{S_L} \sum_{S_R} \int_0^\Delta (P^F + P^A) d\Delta = 1$ for any value of Δ_0 follows $\gamma = \beta - 2\alpha$. Furthermore, if $Q^{F,A}$ describe fixed-point distributions the average gap $\langle \Delta \rangle$ has to scale linearly in Δ_0 , i.e., $\beta = 1$. Finally we argue that $\delta = 2\alpha$. An effective spin at energy scale Δ_0 is built up by two more ‘primitive’ spins which in turn are built up by two even more ‘primitive’ spins etc. The total spin of such a unit is just the sum of the participating original spins, where each spin enters the sum with the same (opposite) sign as its neighbor if the coupling is ferromagnetic (antiferromagnetic). Thus an effective spin made up of n original spins is a sum of n independent random variables and scales as $S^2 \propto n$ (see below). Thus, we find $n \propto S^2 \propto \Delta_0^{-2\alpha}$ and hence $\delta = 2\alpha$.

We have numerically performed the RG for five chains with different starting distributions, including one purely AF distribution and one distribution simulating the low-energy physics of $\text{Sr}_3\text{CuPt}_{0.8}\text{Ir}_{0.2}\text{O}_6$ [10]. In each case we keep the number of sites fixed to 10^6 by adding one link after each decimation and iterate up to $2 \cdot 10^7$ steps. Details of this calculation will be presented elsewhere [11]. In all five chains the distribution after appropriate scaling eventually approaches the universal fixed-point distribution (3). Integrating Q^F and Q^A over two variables gives the distribution of spins and the distribution of gaps plotted in Fig. 1. The ratio of FM bonds to AF bonds is 0.59 at the fixed point and from the scaling of the averages (see Fig. 2) the exponents are determined to be $\alpha = 0.22 \pm 0.01$, $\beta = 1.00 \pm 0.005$ (in both P^F and P^A) and $\delta = 0.44 \pm 0.02$. Thus the scaling form (3) and the relations between the exponents are indeed verified in our numerical calculations, and the fixed-point distributions become

$$P^{F,A} = \frac{1}{\Delta_0^{1-2\alpha}} Q^{F,A}(\Delta/\Delta_0, S_L \Delta_0^\alpha, S_R \Delta_0^\alpha) \quad (4)$$

and

$$n \sim \Delta_0^{-2\alpha} ; \quad \alpha = 0.22 \pm 0.01. \quad (5)$$

The scaling forms (4) and (5) are the key results from which we obtain the magnetic susceptibility χ , entropy σ , and specific heat C . At finite temperature T the RG flows will stop at $\Delta_0 \sim k_B T$. All pairs of spins in links with the gap larger than Δ_0 are frozen to form effective large spins according to the RG, while the spins that have survived down to Δ_0 are typically much more weakly coupled and essentially free. These free spins give a Curie contribution to the magnetic susceptibility per unit length,

$$\frac{\chi}{L} = \frac{\mu^2}{3k_B T} \frac{\langle \mathbf{S}^2 \rangle}{n} = \frac{c}{T} \frac{\Delta_0^{-2\alpha}}{\Delta_0^{-2\alpha}} = \frac{c}{T}. \quad (6)$$

The Curie constant c can be calculated in the following way. Our Hamiltonian (1) contains only nearest-neighbor coupling and, therefore, does not give rise to any frustration. In order to estimate $\langle \mathbf{S}^2 \rangle$, the average size of the effective spins, we consider an isolated system with n spins and search for the total spin quantum number of its ground state. From the multiplet of the ground state we choose the state with the largest z -component, $S_{\text{tot}}^z = S_{\text{tot}}$ (total spin size). This state contains the basis state corresponding to the classical ground state with all spin axes parallel to the z -axis. The correlation of classical spins is given by $S_{\text{cl},i}^z S_{\text{cl},j}^z = S_i S_j \prod_{k=j}^{i-1} \text{sgn}(-J_k)$, where S_i is the size of the spin i . The disorder averaged value of S_{tot}^z can now be obtained within the random walk picture as,

$$\langle (S_{\text{tot}}^z)^2 \rangle = \sum_{i=0}^n (S_{\text{cl},i}^z)^2 + 2 \sum_{i>j} S_{\text{cl},i}^z S_{\text{cl},j}^z$$

$$= n\langle S_i^2 \rangle + 2\langle S_i \rangle^2 \sum_{i=1}^n \sum_{j=0}^{n-1} \prod_{k=j}^{i-1} \text{sgn}(-J_k), \quad (7)$$

where $\langle S_i \rangle$ and $\langle S_i^2 \rangle$ are averages of the spin size of the original spin system (1). Using the average $\langle \text{sgn}(-J_l) \rangle = 2q - 1$ (q : probability that a bond is ferromagnetic) we find the Curie constant,

$$c = \frac{\mu^2}{3k_B} \left[\langle S_i^2 \rangle + \frac{2q-1}{1-q} \langle S_i \rangle^2 \right], \quad (8)$$

in the limit $T \rightarrow 0$ ($n \rightarrow \infty$). In particular, in the case where all spins are $1/2$, the low-temperature Curie constant is $c = (\mu^2/12k_B)q/(1-q)$, which is in contrast to the high-temperature value $c = \mu^2/4k_B$.

The entropy at zero external magnetic field follows from similar arguments. At a given temperature T the spins that have not yet frozen into large spins are essentially free and contribute to the entropy as

$$\frac{\sigma(T, H=0)}{L} \propto \frac{k_B \ln(2\langle S \rangle + 1)}{n} \propto \alpha k_B T^{2\alpha} |\ln T|, \quad (9)$$

which give the leading term of the specific heat as

$$\frac{C(T)}{L} \propto T^{2\alpha} |\ln T|. \quad (10)$$

In a finite magnetic field H and at finite temperature the RG flows are interrupted either by the thermal energy $k_B T$ or by the magnetic energy $E_{\text{ZM}} = \mu \langle S \rangle H$. In the former case the magnetization is given by (6) and the entropy by (9). Otherwise, the magnetic field eventually drives the system away from the fixed point of zero magnetic field into a state of aligned effective spins where the entropy is zero and the magnetization saturates. A magnetic field H starts to align the spins at an energy scale $\Delta_0 \sim \mu \langle S \rangle H$, i.e., $\Delta_0 \sim H^{1/(1+\alpha)}$ and the saturated magnetization can be estimated to be $M/L \sim \mu \langle S \rangle / n \sim H^{\alpha/(1+\alpha)}$. The condition that the chain is not yet dominated by thermal fluctuations thus becomes $k_B T < \Delta_0 \sim H^{1/(1+\alpha)}$ and hence

$$\frac{M(T, H)}{L} \sim \begin{cases} H^{\frac{\alpha}{1+\alpha}}, & T^{1+\alpha} \ll bH \\ H/T, & T^{1+\alpha} \gg bH, \end{cases} \quad (11)$$

where b is a non-universal dimensionful constant. Similarly the entropy $\sigma(T, H)$ decreases rapidly towards zero at $T^{1+\alpha} \sim \mu \langle S_i \rangle H$ when the magnetic field starts to align the spins.

In the random bond AF spin- $\frac{1}{2}$ chain the DM RG treatment becomes asymptotically exact [8] because the perturbation parameter $\varepsilon = \Delta_t/\Delta_0$ approaches zero in the low-energy limit (Δ_t : the typical gap in the chain). This is not the case in our RG scheme where $\varepsilon \sim 0.2$ for the fixed-point distribution (see Fig. 1) and higher order terms induce interactions other than nearest-neighbor

coupling. However, such interactions between spins separated by a distance d have couplings of order ε^d . This exponential dependence on d is preserved in the RG so that the application of an effective nearest-neighbor Hamiltonian (1) is justified. Furthermore, terms of the type $(\mathbf{S}_i \cdot \mathbf{S}_{i+1})^m$ may appear in higher order perturbation. Although they may change the local correlations, they would not affect the relations between the exponents. For this conclusion, it is important to remember that the effective spin scales as $S \propto n^{1/2}$. This scaling results from the random correlation of the spins within clusters of length n and can be understood immediately within a random walk picture, which is quite robust against changes of details of the correlation. A consequence of this is the relation between the exponents, $\alpha = 2\delta$ which reduces the number of free exponents to one, α . The only possible effect of higher order terms is to modify slightly the exponent α . In addition, we emphasize that the Curie-like behavior of the magnetic susceptibility (6) is independent of α .

The low-energy physics of our system is very different from that of the random AF spin- $\frac{1}{2}$ Heisenberg chain studied in Refs. [7–9]. In the AF spin- $\frac{1}{2}$ chain the spins successively lock into singlets which afterwards do not participate in the low-energy physics. This *random singlet phase* [12] is characterized by the formation of singlets between spins that may be far apart from each other. The coupling between two $S = \frac{1}{2}$ spins that have survived down to some low-energy scale is mediated by virtually exciting all the singlets in between the two spins into triplets. This leads to a coupling that decreases exponentially with length, $J = \Delta \propto \exp(-\sqrt{n})$. This, together with the fact that the spin size remains constant ($S = \frac{1}{2}$), leads to susceptibility $\chi \propto T^{-1} |\ln T|^{-2}$ and entropy $\sigma(T, H=0) \propto |\ln T|^{-2}$. In our terminology this would correspond to $P^F = 0$, $\alpha = 0$, and $\delta = 0$ up to logarithmic corrections. In contrast, our fixed point is characterized by large spins and a power law relation between energy and length leading to (5), (6), (9), and (10). It is clear that the two systems belong to different universality classes. We emphasize here that the universality class of our model includes a large variety of random spin models. In particular, an AF spin chain with random spin sizes belongs to this class. (It is easy to see that such a system generates effective FM interaction among effective spins as the RG iteration proceeds.) Similarly, any spin chain with randomly distributed AF and FM bonds should belong to this universality class [13]. Hence the $S = \frac{1}{2}$ AF fixed point studied in Refs. [7–9] is unstable toward the presence of FM couplings as well as toward randomness in the magnitude of the spins. In either case we expect the system to flow to the fixed point discussed in this paper.

An important candidate for this universality class is represented by the compound $\text{Sr}_3\text{CuPt}_{1-x}\text{Ir}_x\text{O}_6$ mentioned earlier, where the low-energy physics is described

by the Heisenberg model (1). This system shows three Curie regimes with different Curie constants [6]. At high temperature it corresponds to independent $S=1/2$ spins, the intermediate regime is governed by free effective spins formed on the FM and AF segments, and the low-temperature regime scales to our universality class. The slow approach of $\langle n \rangle$ to the fixed point shown in Fig. 2 means that it will be difficult to observe the fixed-point value of the specific heat exponent whereas we find numerically that the limit of χ given in (6) and (8) should be more readily observable.

We would like to thank T. K. Ng for helpful discussions and H.-C. zur Loye and T. N. Nguyen for stimulating our interest in the problem. We are also grateful for the financial support by the Swedish Natural Science Research Council (E.W.), by Swiss Nationalfonds (M.S., No. 8220-037229), by the M.I.T. Science Partnership Fund, and by the NSF through the Material Research Laboratory (DMR-90-22933).

* On leave from Department of Applied Physics, University of Tokyo, Hongo, Tokyo 113, Japan.

- [1] A. P. Wilkinson, A. K. Cheetham, W. Kunman, and A. Kvik, *Eur. J. Solid State Inorg. Chem.* **28**, 453 (1991).
- [2] Strictly speaking it is the occurrence of Ir atoms that is random so that FM bonds always appear in pairs.
- [3] T. N. Nguyen, Ph. D thesis, Massachusetts Institute of Technology, 1994 (unpublished); T. N. Nguyen, D. M. Giaquinta, and H.-C. zur Loye, *Chem. Mater.* **6**, 1642 (1994).
- [4] A. Furusaki, M. Sigrist, P. A. Lee, K. Tanaka, and N. Nagaosa, *Phys. Rev. Lett.* **73**, 2622 (1994).
- [5] While a crossover between different Curie regimes also appears in the corresponding model with classical spins, the entropy of the classical model is not sensitive to the disorder and it is clearly necessary to study the quantum

problem.

- [6] A. Furusaki, M. Sigrist, E. Westerberg, P. A. Lee, K. B. Tanaka, and N. Nagaosa, unpublished.
- [7] C. Dasgupta and S.-k. Ma, *Phys. Rev. B* **22**, 1305 (1980); S.-k. Ma, C. Dasgupta, and C. K. Hu, *Phys. Rev. Lett.* **43**, 1434 (1979).
- [8] D. S. Fisher, *Phys. Rev. B* **50**, 3799 (1994), and references therein.
- [9] Hirsch and José studied the random AF spin- $\frac{1}{2}$ Heisenberg chain using Kadanoff's block spin RG scheme, J. E. Hirsch and J. V. José, *Phys. Rev. B* **22**, 5339 (1980); J. E. Hirsch, *ibid.* **22**, 5355 (1980).
- [10] To simulate $\text{Sr}_3\text{CuPt}_{0.8}\text{Ir}_{0.2}\text{O}_6$ we start with a spin- $\frac{1}{2}$ chain with 20% FM and 80% AF bonds (all with strength J) and perform the segmentation described in [6]. This gives the distribution of couplings between the effective degrees of freedom which we take as initial distribution in the DM RG.
- [11] E. Westerberg, A. Furusaki, M. Sigrist, and P. A. Lee, in preparation.
- [12] R. N. Bhatt and P. A. Lee, *Phys. Rev. Lett.* **48**, 344 (1982).
- [13] Obviously random FM systems of any type are excluded from this class.

FIG. 1. (a) The fixed-point distributions of the spin sizes, $P_S^{F,A}(S) \equiv \sum_{S_L} \int d\Delta Q^{F,A}$. In the figure both distributions are normalized so that $\sum_S P_S^{F,A} = 1$. (b) The fixed-point distributions for the gaps, $P_\Delta^{F,A}(\Delta) \equiv \sum_{S_L, S_R} Q^{F,A}$. The distributions are normalized as in (a). Inset: Schematic picture of the decimation procedure.

FIG. 2. Scaling of the averages which determine the exponents α , β , and δ . The averages of the gap $\langle \Delta \rangle$ and spin size $\langle S \rangle$ are calculated from $P^F(\Delta_0; \Delta, S_L, S_R)$. The corresponding averages for P^A are almost identical. The initial distribution simulates $\text{Sr}_3\text{CuPt}_{0.8}\text{Ir}_{0.2}\text{O}_6$ [10]. We have found more rapid crossover to the fixed point for other initial distributions, *e.g.* the flat distribution $P^{F,A} = \text{const.}$, $-1 < J < 1$, $S_{L,R} \in \{1/2, 1, \dots, 3\}$.